

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

C. R. Geoscience ●●● (●●●●) ●●●-●●●



COMPTES RENDUS

GEOSCIENCE

<http://france.elsevier.com/direct/CRAS2A/>

External Geophysics, Climate and Environment

Coupled modeling of global carbon cycle and climate in the Neoproterozoic: links between Rodinia breakup and major glaciations

Yves Godd ris^{a,*}, Yannick Donnadi u^b, C line Dessert^c, Bernard Dupr ^a,
Fr d ric Fluteau^d, Louis M. Fran ois^e, Joseph Meert^f, Anne N d lec^a, Gilles Ramstein^b

^a LMTG, Observatoire Midi-Pyr n es, CNRS/universit  Paul-Sabatier, route de Narbonne, 31062 Toulouse cedex, France

^b LSCE, CNRS/CEA, B t. 701, Orme des Merisiers, 91191 Gif-sur-Yvette cedex, France

^c Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EQ, UK

^d IPGP, tour 24, BP 89, 4, place Jussieu, 75252 Paris cedex 05, France

^e LPAP, universit  de Li ge, 5, avenue de Cointe, B4000 Li ge, Belgium

^f Department of Geological Sciences, University of Florida, 241 Williamson Hall, P.O. Box 112120, Gainesville, FL 32611, USA

Received 23 November 2004; accepted after revision 14 December 2005

Written on invitation of the Editorial Board

Abstract

A coupled climate–geochemical model of new generation (GEOCLIM) is used to investigate the possible causes of the initiation of snowball glaciations during Neoproterozoic times. This model allows the calculation of the partial pressure of atmospheric CO₂ simultaneously with the climate at the continental surface with a rough 2D spatial resolution (10° lat. × 50° long.). We calculate that the breakup of the Rodinia supercontinent, starting 800 Myr ago, results in a global climatic cooling of about 8 °C triggered by enhanced consumption of atmospheric CO₂ resulting from increased runoff over continental surfaces. This increase in runoff is driven by the opening of oceanic basins resulting in an increase of soil moisture sources close to continental masses. This climatic effect of the supercontinent breakup is particularly strong within the 800–700 Ma interval since all continents are located in the equatorial area, where temperature and runoff conditions optimize the consumption of CO₂ through weathering processes. However, this effect alone is insufficient to trigger snowball. We propose that the efficient weathering of fresh basaltic surfaces that erupted during the Rodinia breakup, and were transported to the humid equatorial area through continental plate motion, contributed the necessary CO₂ sink that triggered the ca. 730-Ma Sturtian glacial event. Simulations of the GEOCLIM model for the ca 580-Ma Gaskiers ice age, where all continents are centered on the South Pole, shows that no snowball glaciation can be initiated. The calculated CO₂ partial pressure remains above 1000 ppmv, while a threshold of less than 80 ppmv is required to initiate a snowball glaciation. At that time, a polar configuration does not allow the onset of total glaciation. Nevertheless, a regional glaciation is simulated by the GEOCLIM when the climatic and geochemical (i.e. weathering related) effects of the Pan-African orogeny (~600 Ma) are taken into account. Finally, the question of the role of the paleogeographic setting in the Marinoan snowball event (~635 Ma) is still an open question, since no reliable Marinoan paleogeographic reconstruction exists due to the paucity of paleomagnetic data. **To cite this article:** Y. Godd ris *et al.*, *C. R. Geoscience* ●●● (●●●●).

  2005 Acad mie des sciences. Published by Elsevier SAS. All rights reserved.

* Corresponding author.

E-mail address: godderis@lmtg.obs-mip.fr (Y. Godd ris).

R sum 

Mod lisation coupl e du cycle du carbone et du climat au N oproterozo ique : liens entre la dislocation du supercontinent Rodinia et les glaciations majeures. Un mod le coupl  g ochimie–climat de nouvelle g n ration (GEOCLIM) est utilis  afin d'explorer les causes de l'initiation de glaciations de type « boule de neige »   la fin du Prot rozo ique. Ce mod le permet le calcul de la teneur en CO₂ dans l'atmosph re et du climat de mani re simultan e, avec une r solution spatiale de 10  en latitude sur 50  en longitude. Sur la base de simulations par ce mod le, nous calculons que la dislocation du supercontinent Rodinia (qui commence vers 800 Ma) provoque un refroidissement global du climat de 8  C, cons cutif   l'augmentation de la consommation de CO₂ atmosph rique, elle-m me li e   l'augmentation du ruissellement continental. Cet accroissement du ruissellement est li    l'ouverture de bassins oc aniques, agissant comme autant de sources d'humidit    proximit  des continents. Cet effet climatique de la dislocation d'un supercontinent est particuli rement efficace aux alentours de 800–700 Ma, puisque tous les continents se situent dans la zone  quatoriale, o  les conditions de temp rature et de ruissellement optimisent la consommation de CO₂ par les processus d'alt ration. Quand tous les effets li s   la dislocation de la Rodinia sont pris en compte, y compris l'alt ration tr s rapide des surfaces basaltiques fra ches qui se sont  panch es   la surface des continents durant les phases initiales de la dislocation et qui sont emmen es dans la zone  quatoriale humide par la d rive des continents, une glaciation « boule de neige » est initi e, correspondant   l' v nement glaciaire sturtien (~730 Ma). Des simulations du mod le GEOCLIM de la glaciation Gaskiers (580 Ma), o  les continents sont regroup s aux alentours du p le Sud, montrent qu'une glaciation de type « boule de neige » ne peut  tre initi e. La pression partielle de CO₂ atmosph rique calcul e par GEOCLIM est en effet sup rieure   1000 ppmv, alors que le seuil requis pour initier une glaciation « boule de neige » Gaskiers est de l'ordre de 80 ppmv. Nous en d duisons qu'une configuration polaire des continents n'autorise pas la formation d'une Terre « boule de neige ». N anmoins, une glaciation r gionale est simul e par GEOCLIM   580 Ma, lorsque les effets g ochimiques (impact sur l'alt ration) et climatiques de l'orog ne panafricain sont pris en compte, en accord avec les donn es existantes. Enfin, le r le de la pal og ographie dans le cadre de la glaciation « boule de neige » du Marinoen (~635 Ma) n'a pu  tre test , par manque de donn es pal omagn tiques fiables, ce qui n'autorise pas une reconstruction pal og ographique pr cise. *Pour citer cet article : Y. Godd ris et al., C. R. Geoscience ●●● (●●●).*

  2005 Acad mie des sciences. Published by Elsevier SAS. All rights reserved.

Keywords: Numerical modeling; Carbon cycle; Climates; Neoproterozoic; Rodinia carbon

Mots-cl s : Mod lisation num rique ; Cycle du carbone ; N oproterozo ique ; Rodinia

Version fran aise abr g e

Le N oproterozo ique est une p riode marqu e par des  pisodes de glaciation majeurs [17,20,21,24]. Trois  v nements ont  t  jusqu'  pr sent identifi s : la glaciation sturtienne (vers 750–710 Ma) [10], la glaciation marinoenne (autour de 635 Ma) [21] et la glaciation varangienne ou Gaskiers (580 Ma) [4,45]. Parmi ces trois glaciations, les deux premi res pourraient correspondre   des glaciations totales dites « boule de neige », cons cutives, en particulier,   l'abondance de d p ts glaciaires   basse latitude [10,20,21,25]. Si l'existence de glaciations « boule de neige » durant le N oproterozo ique tend aujourd'hui   faire l'unanimit , les causes de ces  v nements restent largement inexplor es.

Dans le cadre du programme ECLIPSE du CNRS *Comprendre et mod liser les glaciations du N oproterozo ique*, nous avons d velopp  un outil num rique nouveau (GEOCLIM [7]), couplant un mod le climatique de basse r solution (le mod le CLIMBER, 50  de longitude \times 10  de latitude [39]), et un mod le g ochimique d crivant, entres autres, les cycles globaux du carbone et de l'alcalinit  (COMBINE [14]). GEOCLIM permet donc de calculer la teneur en CO₂ atmosph -

rique   l' quilibre, avec une configuration pal og ographique donn e (Fig. 1), et pour un d gazage de la Terre solide fix  (Fig. 2). En effet, la r solution spatiale 2D de GEOCLIM autorise le calcul de la consommation de CO₂ atmosph rique par alt ration des silicates   la surface des continents, avec une r solution spatiale non nulle. Cette r solution permet d' tudier, de mani re explicite, l'impact de la configuration continentale sur le ruissellement continental, premier facteur contr lant l'alt ration des silicates [6,37], ce qui  tait inaccessible aux mod les 0D existants [2]. GEOCLIM nous a permis d'explorer pour la premi re fois l'impact g ochimique et climatique du fractionnement du supercontinent Rodinia, qui d bute aux alentours de 800 Ma [32].

Vers 800 Ma, la Rodinia se trouve localis e   l' quateur et sa dislocation disperse les blocs continentaux le long de celui-ci. Les pal olatitudes maximales atteintes par ces blocs continentaux aux alentours de 730 Ma restent comprises entre 40 N et 40 S [31,32]. Cette configuration continentale est particuli re au N oproterozo ique, et ne se rencontre jamais au cours du Phan rozo ique (Fig. 1).

Une premi re simulation de GEOCLIM   800 Ma calcule une pression partielle de CO₂ de 1830 ppmv,

correspondant   une situation o  l'ensemble de la surface de la Rodinia est libre de glace (Fig. 3A). En revanche, une simulation r alis e   730 Ma, dans une configuration pal og ographique disloqu e le long de l' quateur (Fig. 1), fournit une teneur en CO₂ de 510 ppmv seulement (temp rature moyenne mondiale correspondante de 2  C, Fig. 3B). Cette baisse de la teneur en CO₂ correspond   un refroidissement global majeur de 8  C. Ce refroidissement est li    l'augmentation du ruissellement continental li    l'accroissement de sources d'humidit    proximit  des continents (Fig. 4), elle-m me due   l'ouverture de nombreux bras de mer (dont le proto-Pacifique) [7]. Il se produit   d gazage de la Terre solide constant. C'est la premi re fois que l'effet climatique   long terme de la dislocation d'un supercontinent est estim , en couplage avec les changements dans l'alt ration continentale (Fig. 5).

De plus, la dislocation de la Rodinia est pr c d e et accompagn e de la mise en place de nombreuses provinces basaltiques entre 825 et 725 Ma [15,27,28]. L'alt ration chimique de ces provinces accro t la consommation de CO₂ atmosph rique et aggrave le refroidissement climatique global [5,15]. Cet effet devient particuli rement efficace lorsque ces provinces sont amen es dans la zone  quatoriale par la d rive des plaques continentales, o  le ruissellement et la temp rature  lev es optimisent la consommation de CO₂. C'est particuli rement le cas pour l'une des plus grandes d'entre elles, la province magmatique Laurentienne, d' ge 780 Ma, dont l' ruption se produit vers 30  de latitude, et qui atteint la zone  quatoriale vers 730 Ma, ce qui accro t la consommation de CO₂ atmosph rique (Fig. 5) [15]. La simulation GEOCLIM de l'effet d'augmentation de ruissellement due   la dislocation d'un supercontinent, combin    la consommation accrue de CO₂ par les provinces basaltiques, permet de franchir le seuil de CO₂, sous lequel une glaciation de type boule de neige est initi e (250 ppmv) [7] (Fig. 6).

Concernant la glaciation Gaskiers, la configuration continentale est totalement diff rente de la configuration sturtienne, puisque l'ensemble des masses continentales se trouve localis  dans l'h misph re sud, avec le Gondwana ouest (Afrique de l'Ouest et Am rique du Sud) et la Laurentia situ s   proximit  imm diate du p le (Fig. 1) [46]. Une simulation GEOCLIM avec la configuration pal og ographique   580 Ma calcule une teneur en CO₂ atmosph rique   l' quilibre de 1037 ppmv, prenant en compte l'effet d'accroissement de l'alt ration chimique des continents par alt ration m canique accrue dans la zone des orog nes panafricains. N anmoins, quoiqu'une glaciation r gionale soit initi e dans la simulation [9] (en grande partie li e   la

perturbation de la circulation atmosph rique engendr e par la pr sence de l'orog ne, Fig. 7), elle n'atteint pas l'ampleur d'une glaciation de type « boule de neige », le seuil de CO₂ requis pour une telle glaciation  tant de 77 ppmv.

Enfin, l' pisode glaciaire du Marinoen n'a pu  tre simul  par manque de donn es pal og ographiques fiables et par manque de contraintes temporelles fiables. N anmoins, des  tudes r centes semblent indiquer, d'une part, un  ge plus ancien qu'initialement estim  (635 Ma au lieu de 600 Ma [18]) et, d'autre part, une localisation toujours  quatoriale de plusieurs blocs continentaux [30], deux arguments tendant   sugg rer l'applicabilit  de notre mod le pal og ographique des glaciations n oprotozo iques.

En conclusion, nos r sultats sugg rent que les glaciations « boule de neige » se sont produites dans une fen tre temporelle o  la distribution pal og ographique des continents  tait essentiellement  quatoriale et dispers e. Lorsque la d rive des continents a modifi  cette configuration (vers 580 Ma, la plupart des continents sont dans l'h misph re sud), le temps des glaciations « boule de neige »  tait termin .

1. Introduction

The climate of the Late Proterozoic has received much attention in the recent years, since the suggestion of the occurrence of global glaciations in the Sturtian (730 Ma) and Marinoan (635 Ma) time periods (the so-called 'snowball' glaciations) [17,18,21,24]. A third event (the Gaskiers or Varanger glaciation, 580 Ma) has also been recognized as a major glaciation although probably not a snowball event [25]. Apart from the intense debate regarding the exact amplitude of the glacial events themselves [11,21–23], there is little doubt that these glacial episodes were intense compared to the Pleistocene glaciations, with glaciers reaching sea level near the equator [10]. The snowball hypothesis is probably the most attractive model up to now [21], since it successfully explains most of the extraordinary geological observations regarding the glaciations [20]. However, one of the most critical questions remains the cause of the extreme climate deterioration, which has not been experienced since the Neoproterozoic era.

Among the plausible explanations invoked to explain snowball-like glaciations lies the fainter sun at this time period. However, as a direct and obvious consequence of the negative feedback existing between consumption of atmospheric CO₂ through continental silicate weathering and climate [47], a fainter sun would be totally compensated for by a higher partial pressure of CO₂

into the atmosphere [44,47]. High obliquity ($>54^\circ$) has been also invoked as a plausible cause for unusual location of the glacial deposits, resulting in a colder climate at the equator compared to the Polar Regions [48]. Indeed this hypothesis does not require the onset of a complete snowball glaciation to explain the presence of low-latitude glacial deposits. However, this hypothesis has been tested through climate simulations [8] showing that high obliquity fails to trigger extensive ice-sheets on polar supercontinents. This is a crucial point, since it was the main requirement for the sole proposed mechanism to recover from high to low obliquity in the Williams model [48], invoking climate friction [19]. Furthermore, Levrard and Laskar [26] have dismissed the high-obliquity model by demonstrating that climate friction can only explain a $2\text{--}4^\circ$ decrease of the Earth obliquity over a typical time span of 100 million years, while Williams' model requires a value of 40° decrease.

Another explanation relies on changes in the partial pressure of greenhouse gases. Numerous numerical climate simulations were performed at various partial pressure of atmospheric CO_2 in order to see whether a snowball-like glaciation might be initiated under Neoproterozoic conditions. Note that the answer to this question is still a matter of debate, particularly when coupled ocean-atmosphere models are used [40,41]. Nevertheless, threshold $p\text{CO}_2$ below which runaway ice-albedo feedback drives the Earth into a snowball glaciation ranges from 130 to about 500 ppmv depending on the climate model used. However, all these studies used the partial pressure of atmospheric CO_2 as a boundary condition. This is particularly critical for the 'slushball' model that simulates areas of open ocean along the equator, while all continents are covered by ice down to the equator [22]. This was obtained assuming 700 ppmv of CO_2 in the atmosphere, but this value remains totally arbitrary. Atmospheric CO_2 level is controlled at the million year timescale by the balance between volcanic degassing and silicate weathering consumption [47]. Any hypothesis arguing in favor of a change in $p\text{CO}_2$ should deal with changes in CO_2 degassing, or the strength of the silicate weathering feedback, which is heavily dependent on climatic conditions (temperature and runoff [6,37]).


Methane has been invoked as a possible trigger for snowball events. Pavlov and co-authors [38] suggested that the Neoproterozoic atmosphere might have been methane-rich because of sustained methane degassing from the oxygen-poor ocean. An oxidation event at the end of the Proterozoic might have drastically reduced the methane content of the atmosphere, leading to a rapid decrease in the greenhouse effect and to a

snowball glaciation. However, this scenario may explain the occurrence of only one snowball glaciation. Similarly, an hypothetical release of methane into the atmosphere just prior to the snowball glaciation has been also suggested, that boosted the hydrological cycle on continents resulting in an increase in continental runoff, leading to a decrease in partial pressure of CO_2 consumed by boosted silicate weathering. Once the source of methane is arbitrarily turned down, and if CO_2 has fallen below the required threshold, a snowball Earth might be generated [43]. However, the ad-hoc release of methane, which is required to match the observed Marinoan pre-glacial decrease in marine $\delta^{13}\text{C}$, is difficult to constrain, mainly because the timing of the $\delta^{13}\text{C}$ decrease is not easy to fix. Furthermore, the response of the hydrologic cycle to a sudden burst in the greenhouse effect heavily depends on the paleogeographic configuration (as will be shown below), and might not be assumed as a simple linear response to enhanced air temperature.

Apart from possible but yet to be demonstrated methane degassing prior to the snowball events, the most prominent feature of the global environmental changes is the fragmentation of the long-lived Rodinia supercontinent. This fragmentation started 800 Myr ago [32,33], about 70 million years prior to the first snowball Earth event, and was accompanied by an intense magmatic activity [29]. In this contribution, we summarize the main results obtained within the framework of the Eclipse program pertaining to the global climatic and geochemical impact of supercontinental break-up, and show that it could be the main cause of the triggering mechanism for extreme glaciations in the Neoproterozoic.

2. Testing the impact of Rodinia dislocation on the global carbon cycle and climate: need for new 2D climate-geochemical models

By 800 Ma, the Rodinia supercontinent was probably still assembled or was in the early stages of the break up process [29,31–33] (Fig. 1). Its precise location ca 800 Ma still remains a matter of debate, and Rodinia might have been centered on the equator [32,33], or spread from the equator to the pole [29]. Nevertheless, even in the last configuration, Rodinia might have been driven down to the equator through true polar wander triggered by the initiation of a mantle superplume [29] prior to 750 Ma, followed by the fragmentation of the supercontinent with the spreading of the continental blocks between 40°S and 40°N [31]. In both cases, the fragmentation process can thus be assim-



ilated to the breakup at low latitudes of a supercontinent initially centered on the equator (Fig. 1). The opening of many seaways increased the source of humidity along the borders of the newly formed continental blocks, and most probably precipitations above continents. The subsequent increase in runoff should increase continental weathering, particularly of silicates [5,37], and hence atmospheric CO₂ consumption. Although already suggested in the early 1990s [24] to explain qualitatively snowball glaciations, this scenario has never been quantitatively tested, mainly because numerical models of the long-term carbon cycle and climate were not designed for such experiments. The WHAK [47], BLAG [3], and GEOCARB [2] models are all 0D models, where global air temperature is a simple logarithmic function of pCO₂ and, despite various attempts towards better constraining of fluctuations in runoff as a function of paleogeography [2], runoff is assumed to be a simple linear function of global mean air temperature to the first order. As a result, such models assume that a warmer world is a wetter world. This basic assumption might be probably true for a given stable paleogeography with changing atmospheric partial CO₂ pressure, but cannot be asserted when paleogeography is changing. Furthermore, the impact of the breakup of a supercontinent cannot be simulated with such models, since these geochemical–climate models have no explicit spatial resolution. Calculated partial pressure of atmospheric CO₂ is not sensitive to paleogeography in 0D models. More sophisticated models, such as the Fran ois and Walker model [12] and derived models [13,14], account for the latitudinal dispersion of the continental masses. Furthermore, these global biogeochemical models are fully coupled to a 1D energy balance model, calculating air temperature as a function of latitude, solar constant, latitudinal dispersion of continents, and pCO₂. Although the Fran ois–Walker type models introduce a spatial resolution into long-term geochemical models, they cannot be applied to the Rodinia breakup, for two main reasons: (1) the Rodinia breakup occurred at a roughly constant latitude, all continental blocks moving along the equator [34], which cannot be simulated by a 1D latitudinal model, (2) an

Fig. 1. Continental reconstructions at 800, 750 and 580 Ma [33]. ANS: Arabian Nubian Shield; LIP: large igneous provinces with inferred continental basalt flows (traps); Mad: Madagascar; NC: Napier Complex.

Fig. 1. Pal og ographies   800, 750 et 580 Ma [33]. ANS : Bouclier arabo-nubien ; LIP : grandes provinces magmatiques avec extension probable des coul es basaltiques (traps) ; Mad : Madagascar ; NC : complexe de Napier.




Fig. 2. A schematic description of the GEOCLIM coupled model. The COMBINE model is forced with a given degassing rate, and calculates the sink of carbon through silicate continental weathering with a 2D resolution of 50° long. \times 10° lat. Based on the carbon budget of the oceans and atmosphere boxes, COMBINE calculates the partial pressure of CO_2 in the atmosphere. This $p\text{CO}_2$ is then used by the CLIMBER climate model, which calculates continental air temperature and runoff on the same spatial grid. These new climatic parameters are then used by the COMBINE model that estimates a new coeval $p\text{CO}_2$. GEOCLIM is then run until a steady state is reached for climate and $p\text{CO}_2$ all together.

Fig. 2. Une vue sch matique du mod le coupl  GEOCLIM. Le mod le COMBINE est forc  avec un d gazage fix , et calcule le puits de carbone par alt ration des silicates continentaux avec une r solution spatiale de 50° longitude \times 10° latitude. Sur la base de la r solution du budget de carbone des oc ans et de l'atmosph re, COMBINE calcule la pression partielle du CO_2 dans l'atmosph re. Ce $p\text{CO}_2$ est alors utilis  par le mod le climatique CLIMBER, qui calcule le ruissellement et la temp rature sur la m me grille continentale. Ces nouveaux param tres climatiques sont alors utilis s par COMBINE, qui estime un nouveau $p\text{CO}_2$. Le mod le GEOCLIM tourne ainsi jusqu'  l'obtention d'un  tat stationnaire simultan ment pour le climat et le $p\text{CO}_2$.

energy balance climate model does not calculate explicitly the water cycle, and runoff (and hence continental weathering) cannot be realistically evaluated above continents.

Because of all these limitations, we developed a new numerical model of the global carbon–alkalinity cycles working at the million-year timescale, coupled to a 2D climate model: the GEOCLIM model (Fig. 2). A full description of this model has been previously published [7,9]. The geochemical module consists of a 6-box carbon–alkalinity cycle model (COMBINE, [14]), while the climate module is an Earth model of intermediate complexity (CLIMBER, [39], adapted for Neoproterozoic conditions [8]). Both models are coupled so that COMBINE calculates the partial pressure of atmospheric CO_2 by solving the global carbon budget (including continental silicate weathering, while volcanic degassing is a forcing), and this $p\text{CO}_2$ is used by CLIMBER to calculate continental air temperature and runoff with a resolution of 50° long. \times 10° lat. (details about the coupling procedure can be found in Donnadiu et al. [7]). These climatic variables are used to calculate new continental weathering fluxes (including silicate weathering) with a 2D spatial resolution, and

COMBINE calculates a new $p\text{CO}_2$. GEOCLIM is then run until a steady-state solution for both $p\text{CO}_2$ and climate is found. Steady state is reached when continental silicate weathering balances exactly volcanic CO_2 degassing (requiring about 5 million years simulated).

3. Rodinia dislocation initiates a severe global cooling: the Sturtian glaciation

All simulations were performed assuming a reduction in solar luminosity by 6% relative to the present-day value. The solid Earth CO_2 degassing rate is held constant at its present value (estimated at 6.8×10^{12} mol yr $^{-1}$ in order to balance the present-day CO_2 consumption through silicate weathering [6]), since no reliable constraint on this flux exists for Neoproterozoic times. Although not necessarily realistic in the context of supercontinental breakup, this assumption allows us to isolate the effect of paleogeographic change on climate. Furthermore, while there is no doubt that the Rodinia breakup was accompanied by flood basalt volcanism [15,27,28], there is no evidence for long-term (10^7 years) sustained increased degassing rate related to breakup.

Continental silicate weathering is assumed to be a linear function of continental runoff, and an exponential function of annual mean air temperature with activation energy of $48\,700$ J mol $^{-1}$ [37] for each continental pixel. Since the spatial resolution of the GEOCLIM model is quite crude (50° long. \times 10° lat.), we assume that under equivalent temperature and runoff conditions, the consumption of CO_2 per square meter through silicate weathering is the same for all continental pixels. This assumption is equivalent to the assumption that the ratio between silicate outcrops versus total surface within each continental pixel is constant.

A first simulation was performed using the 800-Ma Rodinia reconstruction of [31] and [33], to calculate steady state $p\text{CO}_2$ prior to continental breakup. GEOCLIM calculated a coeval steady-state $p\text{CO}_2$ of 1830 ppmv [7], a positive annual mean air temperature for all continental surfaces (Fig. 3A), and global mean annual temperature of 10.8°C . Continental surfaces are quite dry, with the northern part of Rodinia being the driest area (Fig. 4). This explains why the partial pressure of CO_2 is so high, while degassing rate is fixed at its present-day value. Dry conditions reduce the efficiency of silicate weathering, hence leading to an increase in $p\text{CO}_2$ and a warming of global climate.

A second GEOCLIM simulation was then performed using the 750 Ma post-breakup continental configuration of [31] and [33]. The steady-state partial pres-




Fig. 3. Annual mean air temperature ($^{\circ}\text{C}$) calculated by the GEOCLIM model for (A) the 800 Ma paleogeographic setting, (B) 750 Ma.

Fig. 3. Temp rature moyenne annuelle ($^{\circ}\text{C}$) calcul e par le mod le GEOCLIM pour (A) la configuration pal og ographique   800 Ma, (B)   750 Ma.




Fig. 4. Continental runoff in cm yr^{-1} as a function of latitude as calculated by the GEOCLIM model at a $p\text{CO}_2$ of 1830 ppmv. Solid line is for the 800-Ma paleogeographic setting, and dashed line for 750 Ma (adapted from [7]).

Fig. 4. Ruissellement continental (cm an^{-1}) en fonction de la latitude, calcul e par le mod le GEOCLIM   1830 ppmv. La ligne continue repr sente la configuration pal og ographique   800 Ma, et la ligne pointill e la configuration   750 Ma (adapt e de [7]).




Fig. 5. Continental silicate weathering (in 10^4 mol of Ca^{2+} or Mg^{2+} $\text{km}^{-2} \text{yr}^{-1}$) calculated by GEOCLIM at (A) 800 Ma, and (B) 750 Ma. The dashed circle locates the Laurentian magmatic province. Adapted from [7].

Fig. 5. Alt ration des silicates continentaux (en 10^4 mol de Ca^{2+} ou Mg^{2+} $\text{km}^{-2} \text{an}^{-1}$), calcul e par le mod le GEOCLIM   (A) 800 Ma, (B) 750 Ma. Le cercle pointill  repr sente la position de la province magmatique Laurentienne. Adapt e de [7].

sure of CO_2 declined to 510 ppmv in this model run, while mean annual air temperature fell below 0°C over large areas of the continents (mainly above 40 to 45° lat.), as previously established by [7] (Fig. 3B). Global mean annual air temperature fell by more than 8°C from the 800-Ma configuration to the 750-Ma configuration (characterized by a mean global air temperature of 2°C), corresponding to a negative radiative forcing of 6.85 W m^{-2} . The reason for this decrease lies in the increase of continental runoff through the opening of seaways in between continental masses, enhancing the supply of humidity to continents, finally resulting in an enhanced runoff and consumption of CO_2 through silicate weathering (Fig. 5). It is noteworthy that solid Earth degassing rate is kept the same for both 800- and 750-Ma simulations, while climate changes radically, a conclusion that cannot be reached

with 0D/1D models for this paleogeographic configuration.

We conclude that the breakup of a supercontinent exerts a strong cooling effect, especially when the breakup occurs at low latitude as it did in the Neoproterozoic. However, despite a strong cooling, this process alone cannot initiate a snowball glaciation, since the partial pressure of CO₂ reached at 750 Ma is approaching, but still above the pCO₂ threshold required to generate the climatic instability leading to a snowball glaciation [20] (here, the CLIMBER 2 model calculates this threshold at 250 ppmv [7], a value compatible with previous studies).

4. Initiating a Sturtian snowball Earth through basaltic surface weathering

As already mentioned above, the breakup of the Rodinia supercontinent is accompanied by the eruption of large magmatic provinces between 825 and 723 Ma [15,27,28]. As already demonstrated for the Deccan traps at the K–T boundary [5], the subsequent weathering of these fresh basaltic surfaces initiates a long term global climate cooling. Global cooling during the Rodinian breakup would be enhanced. As demonstrated in Godd ris et al. [15] and Donnadi u et al. [7], the weathering of the Laurentian magmatic province (erupted at 780 Ma, with a size arbitrarily fixed to 7×10^6 km², a size comparable to the Siberian traps just after their eruption) might drive the Earth into a snowball, by enhancing the global cooling initiated by the increase in runoff triggered by the dislocation itself. This supplemental cooling forcing drove the icecap to reach 30° latitude, where the resulting climatic instability forced the Earth system into a snowball state. The cooling effect (additional negative radiative forcing of 3.8 W m^{-2}) resulting from the weathering of basaltic surfaces is particularly efficient, since the Laurentia magmatic province, initially in the sub-tropic area (the driest simulated region), where chemical weathering is inefficient, is rapidly driven into low-latitude area through continental plate motions around 750–710 Ma, where runoff and temperature conditions optimize the weathering and subsequent consumption of atmospheric CO₂ (Fig. 5).

As a result, we suggest that the breakup of the Rodinia supercontinent starting 800 Myr ago is the main culprit of the onset of the Sturtian snowball glaciation about 50 or more million years later, through enhanced weathering triggered by enhanced runoff, and weathering of fresh basaltic surfaces (Fig. 6).




Fig. 6. Schematic representation of the onset of the Sturtian snowball glaciation. Calculated pCO₂ is on the Y-axis. The blue/grey area represents the CO₂ values at which the Earth is totally frozen (calculated by CLIMBER). The dashed line shows the upper and lower limits of the pCO₂ required to settle a snowball glaciation calculated by a variety of other climate models. Time is running from left to right. Stage 1 corresponds to the calculated pCO₂ for the Rodinia supercontinental configuration, stage 2 is the calculated pCO₂ for the Rodinia supercontinent with the Laurentia magmatic province (first located in dry area, where only slow weathering and CO₂ consumption occur), and stage 3 stands for the 750-Ma dislocated configuration where the Laurentia magmatic province has been driven to the equatorial humid area. Stage 3 corresponds to a snowball glaciation. The total decrease in the radiative forcing from 800 Ma to the snowball onset reaches more than 10 W m^{-2} .

Fig. 6. Repr sentation sch matique de la mise en place d'une glaciation « boule de neige » au Sturtien. La pCO₂ calcul e est repr sent e par l'axe des Y. La surface bleue (ou en gris e) repr sente les valeurs de CO₂ pour lesquelles la Terre est totalement englac e (calcul es par CLIMBER). Les lignes pointill es repr sentent les limites sup rieure et inf rieure pour lesquelles une glaciation « boule de neige » est produite, pour divers autres mod les climatiques. Le temps s' coule de la droite vers la gauche. L' tape 1 correspond   la pCO₂ calcul e pour une configuration supercontinentale (Rodinia). L' tape 2 est la teneur en CO₂ calcul e pour la configuration Rodinia, avec en plus la pr sence de la province magmatique Laurentienne (d'abord localis e dans une zone   faible ruissellement, ce qui limite fortement l'alt ration et la consommation de CO₂). L' tape 3 correspond   la configuration pal og ographique disloqu e   750 Ma, o  la province magmatique Laurentienne a atteint la zone  quatoriale humide. L' tape 3 correspond   une Terre « boule de neige ». La baisse totale du for age radiatif de 800 Ma   la mise en place de la glaciation boule de neige d passe 10 W m^{-2} .

5. Simulating the Gaskiers glaciation (580 Ma): no snowball

A second set of GEOCLIM simulations was performed using the 580-Ma paleogeographic reconstruction [9], which corresponds to the Gaskiers glacial deposit in Newfoundland [4]. This glaciation was first correlated with the Marinoan event, which is the strongest candidate for a snowball event. However, recent work indicates that the 580-Ma glaciation post-dates the Marinoan glaciation, and although likely severe, was not a snowball [1,10,25,42], and may have been short-lived [4].




Fig. 7. Mean annual air temperature calculated by the GEOCLIM model for the 580-Ma paleogeographic setting ($p\text{CO}_2 = 1037$ ppmv). Pan-African orogen is located by a grey shaded contour. Perennial snow can accumulate above the West Gondwana (where the presence of the Pan-African orogen results in very low temperatures) and part of the Laurentia cratons. Surface winds are also shown for the latitudes 30°S – 90°S , the size of the vector below the panel is for a wind of 8 m s^{-1} . Adapted from [9].

Fig. 7. Temp erature moyenne annuelle de l'air calcul ee par le mod ele GEOCLIM pour la configuration pal eog eographique   580 Ma ($p\text{CO}_2 = 1037$ ppmv). Les conditions climatiques sont requises pour l'accumulation de neige p erennelle sur le Gondwana ouest (o  la pr esence de l'orog ene panafricain impose des temp eratures tr es basses), et sur une partie de la Laurentia. Les vents en surface sont repr esent es par les fl eches (la fl eche sous le graphique repr esente un vent de 8 m s^{-1}). Adapt e de [9].

The paleogeographic setting at 580 Ma is totally different compared to the setting at 750 Ma [46]. Most of the continental masses are located in the southern hemisphere, with West Gondwana and Laurentia located near the South Pole (Fig. 1). Furthermore, the continental topography is dominated by highlands related to the Pan-African orogeny. Under these paleogeographic conditions, the $p\text{CO}_2$ threshold required to initiate a snowball glaciation was calculated by the CLIMBER climate model at 77 ppmv, while the steady-state $p\text{CO}_2$ calculated by a GEOCLIM simulation stabilizes at 1477 ppmv (with present-day CO_2 degassing), corresponding to a mean global air temperature of 10.8°C [9], far above the threshold for snowball initiation (Fig. 7). This high steady-state $p\text{CO}_2$ is mainly the result of the polar location of the continental masses, where low temperatures inhibit chemical weathering. Going below the 77 ppmv threshold through reduction of the degassing would require unrealistic lowering of solid Earth degassing rates, or unrecognized controls on silicate weathering. One such control might be the impact of high rates of mechanical erosion in the Pan-African mountain belts, which would increase chemical weathering rates by increasing the area available for these reactions to occur. We have tested this effect on

our 580 Ma reconstruction by assuming that mechanical erosion in the Pan-African orogen is roughly three times higher than mechanical erosion in low lands, similar to observed rates in the Himalayan orogen [35]. Based on the work by Millot et al. [36], we multiply the spatialized weathering rates by a factor $f_{\text{mech}}^{0.66}$, with $f_{\text{mech}} = 1$ in flat lands, and 3 inside the Pan-African mountain belts. Despite this additional CO_2 consumption, the steady-state $p\text{CO}_2$ calculated by the GEOCLIM model falls to 1037 ppmv, still far from the snowball threshold.

However, a regional-scale glaciation is generated in any case, with possible development of ice sheets on the Gondwana and Laurentia cratons. In the GEOCLIM simulation, accumulation of perennial snow above these cratons is largely driven by the strong perturbation in the atmospheric circulation induced by the Pan-African mountain belts [9] (Fig. 7) (a GEOCLIM simulation performed assuming flat continents does not show any perennial snow accumulation).

Therefore, GEOCLIM simulations for the Gaskiers ice age suggest the development of a severe regional glaciation [42], but do not allow the initiation of a snowball Earth, which is consistent with geological evidence that does not indicate global glaciation at this time [25].

6. The Marinoan glaciation: a question of paleogeographic setting

The Marinoan event is probably the best documented Neoproterozoic glaciation, and a strong candidate for a snowball Earth event at 635 Ma [18]. However, the lack of well-defined paleomagnetic poles for this time precludes any precise paleogeographic reconstructions, and GEOCLIM simulations cannot be confidently performed. Recent studies suggest that an equatorial location of continental masses might still apply to the Marinoan ice age [30]. The question whether this configuration can be qualified as 'dispersed' is open. If continents were still in a dispersed configuration similar to the Sturtian one, then partial pressure of CO_2 should have been kept at low levels, and the global climate prior to the snowball glaciation should be rather chilly, facilitating the onset of a snowball Earth. Nevertheless, the final trigger leading to complete glaciation was most probably different from the Sturtian to the Marinoan ice age, as recorded in the $\delta^{13}\text{C}$ of marine carbonates just prior to the onset of the snowball: the Sturtian record shows a small increase of the $\delta^{13}\text{C}$, while a strong decrease reaching 10‰ is observed worldwide prior to the Marinoan glaciation [16].

7. Uncertainties and limitations

This study aims at understanding the coupled behavior of the global geochemical cycles and climate during the Neoproterozoic. It integrates numerous processes that are sometimes difficult to constrain, particularly for the distant past, and further investigations are strongly required to quantify and/or constrain each of these processes in order to validate our scenario. Among them lies:

- (a) the solid Earth CO₂ degassing, that was held constant in this study mainly due to the lack of constraints, but is suspected to fluctuate during a supercontinent breakup;
- (b) the paleogeography, particularly for Marinoan times, and the timing of the glacial events;
- (c) the question of the precise extent and location of basaltic provinces that are almost totally eroded now;
- (d) the exact geometry of the Pan-African mountain belts, and clues of their mechanical erosion rate.

Constraining all these uncertainties is a challenge for future research aiming at understanding the catastrophic glaciations that occurred at the end of the Neoproterozoic.

8. Conclusions

The GEOCLIM model simulations suggest that the Sturtian glaciation was triggered by the Rodinia breakup at low latitudes. By the time of the Gaskiers glaciation (580 Ma), a dispersed continental configuration along the equator was no more topical, and continental drift towards the South Pole precluded the initiation of a snowball glaciation. We suggest that such extreme glaciations were greatly facilitated by the dispersal of an equatorial supercontinent that increased continental runoff, enhanced consumption of atmospheric CO₂ through silicate weathering and created the conditions for the initiation of the snowball events. The final trigger might have been different for the Sturtian and Marinoan events. We suggest that weathering of large basaltic provinces erupted during the early stages of the Rodinia breakup was responsible for the Sturtian snowball glaciation, adding its cooling effect on global climate to the runoff enhancement effect of the Rodinia dislocation. Our results suggest that this control on atmospheric CO₂ was for more than 60% of the global cooling leading to the snowball. As a result, the time window for snowball Earth events ranges

from the fragmentation of the Rodinia supercontinent (ca 800 Ma) until the convergence of West Gondwana and Laurentia around the South Pole (ca 580 Ma). An exclusively equatorial paleogeographic configuration has not occurred since, explaining why the Phanerozoic glaciations were never as intense as the Neoproterozoic events.

Acknowledgements

This work was achieved thanks to the financial support of the CNRS through the Eclipse program ‘‘Comprendre et mod liser les glaciations du N oproterozoique’’.

References

- [1] G.H. Barfod, F. Albar de, A.H. Knoll, S. Xiao, et al., New Lu–Hf and Pb–Pb age constraints on the earliest animal fossils, *Earth Planet. Sci. Lett.* 201 (2002) 203–212.
- [2] R.A. Berner, Z. Kothavala, GEOCARB III: A revised model of atmospheric CO₂ over Phanerozoic time, *Am. J. Sci.* 301 (2001) 182–204.
- [3] R.A. Berner, A.C. Lasaga, R.M. Garrels, The carbonate–silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 millions years, *Am. J. Sci.* 284 (1983) 641–683.
- [4] S.A. Bowring, P. Myrow, E. Landing, J. Ramenzani, Geochronological constraints on terminal Neoproterozoic events and the rise of metazoans, in: *NASA Astrobiology Institute General Meeting*, Arizona state University, Tempe, Arizona, 2003, p. 113.
- [5] C. Dessert, B. Dupr , L.-M. Fran ois, J. Schott, et al., Erosion of Deccan Traps determined by river geochemistry: impact on the global climate and the ⁸⁷Sr/⁸⁶Sr ratio of seawater, *Earth Planet. Sci. Lett.* 188 (3–4) (2001) 459–474.
- [6] C. Dessert, B. Dupr , J. Gaillardet, L.M. Fran ois, et al., Basalt weathering laws and the impact of basalt weathering on the global carbon cycle, *Chem. Geol.* 202 (2003) 257–273.
- [7] Y. Donnadi u, Y. Godd ris, G. Ramstein, A. N elec, et al., Snowball Earth triggered by continental breakup through changes in runoff, *Nature* 428 (2004) 303–306.
- [8] Y. Donnadi u, G. Ramstein, F. Fluteau, D. Roche, et al., The impact of atmospheric and oceanic heat transport on the sea-ice-albedo instability during the Neoproterozoic, *Clim. Dynam.* 22 (2–3) (2004) 293–306.
- [9] Y. Donnadi u, G. Ramstein, Y. Godd ris, F. Fluteau, Global tectonic setting and climate of the Late Neoproterozoic: a climate-geochemical coupled study, in: G. Jenkins, M. McMenamin, L. Sohl, C. McKay (Eds.), *The Extreme Proterozoic: Geology, Geochemistry, and Climate*, in: *Geophys. Monogr.*, vol. 146, 2004, p. 200.
- [10] D.A.D. Evans, Stratigraphic, geochronological, and paleomagnetic constraints upon the Neoproterozoic climatic paradox, *Am. J. Sci.* 300 (2000) 347–433.
- [11] N. Eyles, N. Januszczak, ‘Zipper-rift’: a tectonic model for Neoproterozoic glaciations during the breakup of Rodinia after 750 Ma, *Earth Sci. Rev.* 65 (2004) 1–73.
- [12] L.M. Fran ois, J.C.G. Walker, B.N. Opdyke, The history of global weathering and the chemical evolution of the ocean–atmosphere system, in: E. Takahashi, R. Jeanloz, D. Dubie

- (Eds.), *Evolution of the Earth and Planets*, vol. 14, International Union of Geodesy and Geophysics and the American Geophysical Union, Washington, 1993, pp. 143–159.
- [13] Y. Godd ris, L.-M. Fran ois, The Cenozoic evolution of the strontium and carbon cycles: relative importance of continental erosion and mantle exchanges, *Chem. Geol.* 126 (1995) 169–190.
- [14] Y. Godd ris, M.M. Joachimski, Global change in the Late Devonian: modelling the Frasnian–Famennian short-term carbon isotope excursions, *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 202 (2004) 309–329.
- [15] Y. Godd ris, A. N d lec, Y. Donnadiou, B. Dupr , et al., The Sturtian glaciation: Fire and ice, *Earth Planet. Sci. Lett.* 211 (2003) 1–12.
- [16] G.P. Halverson, A.C. Maloof, P.F. Hoffman, The Marinoan glaciation (Neoproterozoic) in Northeast Svalbard, *Basin Res.* 16 (2004) 297–324.
- [17] W.B. Harland, M.S. Rudwick, The great infra-Cambrian ice age, *Sci. Am.* 211 (1964) 28–36.
- [18] K.-H. Hoffmann, D.J. Condon, S.A. Bowring, J.L. Crowley, U–Pb zircon date from the Neoproterozoic Ghaub formation, Namibia: constraints on Marinoan glaciation, *Geology* 32 (2004) 817–820.
- [19] P.F. Hoffman, A.C. Maloof, Glaciation: the snowball theory still holds water, *Nature* 397 (1999) 384.
- [20] P.F. Hoffman, D.P. Schrag, The snowball Earth hypothesis: testing the limits of global change, *Terra Nova* 14 (2002) 129–155.
- [21] P.F. Hoffman, A.J. Kaufman, G.P. Halverson, D.P. Schrag, A Neoproterozoic Snowball Earth, *Science* 281 (1998) 1342–1346.
- [22] W.T. Hyde, T.J. Crowley, S.K. Baum, W.R. Peltier, Neoproterozoic ‘snowball Earth’ simulations with a coupled climate/ice sheet model, *Nature* 405 (2000) 425–429.
- [23] M.J. Kennedy, N. Christie-Blick, A.R. Prave, Carbon isotopic composition of Neoproterozoic glacial carbonates as a test of paleoceanographic models for snowball Earth phenomena, *Geology* 29 (2001) 1135–1138.
- [24] J.L. Kirschvink, Late Proterozoic low-latitude global glaciation: the snowball Earth, in: J.W. Schopf, C. Klein (Eds.), *The Proterozoic Biosphere*, Cambridge University Press, Cambridge, 1992, pp. 51–52.
- [25] A.H. Knoll, M.R. Walter, G.M. Narbonne, N. Christie-Blick, A new period for the geologic time scale, *Science* 305 (2004) 621–622.
- [26] B. Levrard, J. Laskar, Climate friction and the Earth’s obliquity, *Geophys. J. Inter.* 154 (2003) 970–990.
- [27] Z.X. Li, X.H. Li, P.D. Kinny, J. Wang, The break up of Rodinia: did it start with a mantle plume beneath South China?, *Earth Planet. Sci. Lett.* 173 (1999) 171–181.
- [28] Z.X. Li, X.H. Li, P.D. Kinny, J. Wang, et al., Geochronology of Neoproterozoic syn-rift magmatism in the Yangtze Craton, South China and correlations with other continents: evidence for a mantle superplume that broke up Rodinia, *Precambrian Research* 2286 (2003) 1–25.
- [29] Z.X. Li, D.A.D. Evans, S. Zhang, A 90° spin on Rodinia: possible causal links between the Neoproterozoic supercontinent, superplume, true polar wander and low-latitude glaciation, *Earth Planet. Sci. Lett.* 220 (2004) 409–421.
- [30] M. Macouin, J. Besse, M. Ader, S. Gilder, et al., Combined paleomagnetic and isotopic data from the Doushantuo carbonates, South China: implications for the ‘snowball Earth’ hypothesis, *Earth Planet. Sci. Lett.* 224 (2004) 387–398.
- [31] J.G. Meert, A synopsis of events related to the assembly of eastern Gondwana, *Tectonophysics* 362 (2003) 1–40.
- [32] J.G. Meert, C.M. Powell, Assembly and break-up of Rodinia: introduction to the special volume, *Precamb. Res.* 110 (2001) 1–8.
- [33] J.G. Meert, T.H. Torsvik, The making and unmaking of a supercontinent: Rodinia revisited, *Tectonophysics* 375 (2003) 261–288.
- [34] J.G. Meert, T.H. Torsvik, Paleomagnetic constraints on Neoproterozoic ‘Snowball Earth’ continental reconstructions, in: G.S. Jenkins, M. McMenamin, C.P. McKay, L. Sohl (Eds.), *The Extreme Proterozoic: Geology, Geochemistry, and Climate*, in: *Geophys. Monogr.*, vol. 146, 2004, pp. 5–11.
- [35] F. M tievier, Y. Gaudemer, P. Tapponnier, M. Klein, Mass accumulation rates in Asia during the Cenozoic, *Geophys. J. Int.* 137 (1999) 280–318.
- [36] R. Millot, J. Gaillardet, B. Dupr , C.J. All gre, The global control of silicate weathering rates and the coupling with physical erosion: new insights from rivers of the Canadian Shield, *Earth Planet. Sci. Lett.* 196 (2002) 83–98.
- [37] P. Oliva, J. Viers, B. Dupr , Chemical weathering in granitic crystalline environments, *Chem. Geol.* 202 (2003) 225–256.
- [38] A.A. Pavlov, M.T. Hurtgen, J.F. Kasting, M.A. Arthur, Methane-rich Proterozoic atmosphere, *Geology* 31 (2003) 87–90.
- [39] V. Petoukhov, A. Ganopolski, V. Brovkin, M. Claussen, et al., CLIMBER-2: a climate system model of intermediate complexity. Part I: Model description and performance for present climate, *Clim. Dynam.* 16 (2000) 1–17.
- [40] C.J. Poulsen, R.L. Jacob, R.T. Pierrehumbert, T.T. Huynh, Testing paleogeographic controls on a Neoproterozoic snowball Earth, *Geophys. Res. Lett.* 29 (2002).
- [41] C.J. Poulsen, R.T. Pierrehumbert, R.L. Jacob, Impact of ocean dynamics on the simulation of the Neoproterozoic ‘snowball Earth’, *Geophys. Res. Lett.* 28 (2001) 1575–1578.
- [42] A.H.N. Rice, G.P. Halverson, P.F. Hoffman, Three for the Neoproterozoic: Sturtian, Marinoan and Varangerian glaciations, *Geophys. Res. Abstr.* 5 (2003) 11425.
- [43] D.P. Schrag, R.A. Berner, P.F. Hoffman, G.P. Halverson, On the initiation of a Snowball Earth, *Geochem. Geophys. Geosyst.* 3 (2002), doi:10.1029/2001GC000219.
- [44] E. Tajika, Faint young sun and the carbon cycle: implication for the Proterozoic global glaciations, *Earth Planet. Sci. Lett.* 214 (2003) 443–453.
- [45] M.D. Thomson, S.A. Bowring, Age of the Squantum ‘tillite’, Boston basin, Massachusetts: U–Pb zircon constraints on terminal Neoproterozoic glaciation, *Am. J. Sci.* 300 (2000) 630–655.
- [46] R. Trompette, Gondwana evolution; its assembly at around 600 Ma, *C. R. Acad. Sci., Ser. IIA* 330 (2000) 305–315.
- [47] J.C.G. Walker, P.B. Hays, J.F. Kasting, A negative feedback mechanism for the long-term stabilization of Earth’s surface temperature, *J. Geophys. Res.* 86 (1981) 9776–9782.
- [48] G.E. Williams, Late Precambrian glacial climate and the Earth’s obliquity, *Geol. Mag.* 112 (1975) 441–465.